

$B \rightarrow \rho\pi$ decays, resonant and nonresonant contributions

A. Deandrea^a and A.D. Polosa^b

^a Theory Division, CERN, CH-1211 Genève 23, Switzerland

^b Physics Department, POB 9, FIN-00014, University of Helsinki, Finland

Abstract

We point out that a new contribution to B decays to three pions is relevant in explaining recent data from the CLEO and BABAR collaborations, in particular the results on quasi-two-body decays via a ρ meson. We also discuss the relevance of these contribution to the measurement of CP violations.

PACS: 13.25.Hw, 12.39.Hg
August 2000

$B \rightarrow \rho\pi$ decays, resonant and nonresonant contributions

A. Deandrea^(a) and A.D. Polosa^(b)

^(a) *Theory Division, CERN, CH-1211 Genève 23, Switzerland*

^(b) *Physics Department, POB 9, FIN-00014, University of Helsinki, Finland*
(August, 2000)

We point out that a new contribution to B decays to three pions is relevant in explaining recent data from the CLEO and BABAR collaborations, in particular the results on quasi-two-body decays via a ρ meson. We also discuss the relevance of these contribution to the measurement of CP violations.

Several exclusive charmless hadronic B decays are known with good accuracy, and more will be measured in the near future, due to the large amount of data coming from e^+e^- machines such as the CLEO experiment [1] at the Cornell Electron Storage Ring (CESR), the BaBar experiment at SLAC [2] and the BELLE experiment at KEK [3], or hadron machines such as the Large Hadron Collider (LHC) at CERN, with its program for B -physics. In the following we will consider quasi-two-body B decays to three pions. Our motivation is twofold.

(1) $\bar{B}^0 \rightarrow \rho^\mp \pi^\pm$ and $B^- \rightarrow \rho^0 \pi^-$ were recently measured by the CLEO and BABAR collaborations. The ratio of these branchings are consistent among the two experiments and different from the theoretical expectations.

(2) quasi-two-body B decays to three π can be used for the determination of the angles of the unitarity triangle of CP violations [4].

Concerning the first point, the CLEO collaboration finds [5] $\mathcal{B}(B^- \rightarrow \rho^0 \pi^-) = (10.4^{+3.3}_{-3.4} \pm 2.1) \times 10^{-6}$ and $\mathcal{B}(\bar{B}^0 \rightarrow \rho^\mp \pi^\pm) = (27.6^{+8.4}_{-7.4} \pm 4.2) \times 10^{-6}$, combining these numbers we find a ratio

$$R = \frac{\mathcal{B}(\bar{B}^0 \rightarrow \rho^\mp \pi^\pm)}{\mathcal{B}(B^- \rightarrow \rho^0 \pi^-)} = 2.65 \pm 1.8, \quad (1)$$

where the error in the ratio includes all errors from the branchings; the error may be smaller than what indicated if part of the systematics simplifies in the ratio. The BABAR collaboration [6] gives the following preliminary results $\mathcal{B}(B^- \rightarrow \rho^0 \pi^-) = (24 \pm 8 \pm 3) \times 10^{-6}$ and $\mathcal{B}(\bar{B}^0 \rightarrow \rho^\mp \pi^\pm) = (49 \pm 13^{+6}_{-5}) \times 10^{-6}$, with these numbers we find a ratio $R = 2.0 \pm 1.3$.

As discussed in [7], this ratio is rather small with respect to theoretical expectations; as a matter of fact, when computed in simple approximation schemes, as factorization with no penguins, one gets, from the BSW model [8], $R \simeq 6$. A calculation including penguins in the factorization approximation was performed in [9] and

gives $R \simeq 5.5$. Calculations beyond factorization were performed in [10] with similar results [11] (see Table I).

Previous papers [9,11–13] investigated to role of the B^* and higher resonances in these decays, while here we will investigate a completely different mechanism that enhances selectively some of the $B \rightarrow 3\pi$ decays, namely the possibility that a broad light scalar resonance is present in the 3-body Dalitz plot.

The light σ resonance has accumulated considerable interest after it was reintroduced as a very broad resonance into the 1996 edition of the Reviews of Particle Physics [14]. The E791 collaboration has an evidence of a very broad scalar resonance [15] having mass $m_\sigma = 478 \pm 24$ MeV and width $\Gamma_\sigma = 324 \pm 41$ MeV taking part into the decay process $D^+ \rightarrow 3\pi$ via the resonant channel $D^+ \rightarrow \sigma \pi^+ \rightarrow 3\pi$. The broad σ was controversial due to the extreme difficulty in disentangling it from available data on $\pi\pi \rightarrow \pi\pi$. In the following we will use the the experimental numbers for the mass and width of the σ given by the E791 collaboration. In the process $D^+ \rightarrow 3\pi$ the σ is a good description of three π events not due to quasi-two-body decays of a narrow resonance as it accounts for 46% of the three π 's branching [16]. We will model $B^+ \rightarrow 3\pi$ in a similar way and see that indeed such a contribution is relevant and larger than or comparable to the non-resonant long-distance contribution of the B^* . These two contributions, after the appropriate experimental cuts, constitute an irreducible background to the processes $B \rightarrow \rho\pi$, therefore adding events to some of them and modifying the ratio of branching ratios. This has also consequences on methods based on $B \rightarrow \rho\pi$ for the determination of the CP violating angles. It should be mentioned that the latest edition of the Reviews of Particle Physics [14] gives a broad range for the σ mass and width. Their precise value is not crucial to our analysis as long as the the sigma is within the strip of the Dalitz plot where the ρ mass contribution (within few hundred MeV due to the experimental cuts) is found.

The amplitude we are interested in are those corresponding to the diagrams in Figs. 1 and 2. We compute them at the tree level using the following effective Hamiltonian [8]:

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{ub}^* V_{ud} \{ a_1 (\bar{u}b)_{V-A} (\bar{d}u)_{V-A} + a_2 (\bar{d}b)_{V-A} (\bar{u}u)_{V-A} \}, \quad (2)$$

giving, in the first case:

$$\langle \sigma \pi^- | H_{\text{eff}} | B^- \rangle = \frac{G_F}{\sqrt{2}} V_{ub}^* V_{ud} a_1 F_0^{(B\sigma)} m_\pi^2 (m_B^2 - m_\sigma^2) \times \frac{f_\pi g_{\sigma\pi^+\pi^-}}{\sqrt{2}} \left(\frac{1}{u - m_\sigma^2 + i\Gamma_\sigma(u)m_\sigma} + \frac{1}{t - m_\sigma^2 + i\Gamma_\sigma(t)m_\sigma} \right) \quad (3)$$

where the calculation of the form factor $F_0^{(B\sigma)}$ proceeds as in [17], $g_{\sigma\pi^+\pi^-} = 2.52$ GeV comes from the experimental value of Γ_σ and the co-moving width $\Gamma_\sigma(x)$ is

$$\Gamma_\sigma(x) = \Gamma_\sigma \frac{m_\sigma}{\sqrt{x}} \frac{\sqrt{x/4 - m_\pi^2}}{\sqrt{m_\sigma^2/4 - m_\pi^2}}, \quad (4)$$

where x is $u = (p - p_1)^2$ or $t = (p - p_2)^2$ (as in the crossed channel, see Fig. 1, in which we have two identical π^- particles in the final state). The comoving width can be obtained comparing the usual formula for the fixed width (see for example [14]) with a similar formula where the σ mass is replaced by the square root of the relevant Mandelstam variable (as the σ is an intermediate state in the decay process). The coefficient a_1 is given by $a_1 = C_1 + \frac{1}{3}C_2$ where the Wilson coefficients C_1 and C_2 , fitted for B decays, are $C_1(m_b) = 1.105$ and $C_2(m_b) = -0.228$.

On the other hand, the diagram in Fig. 2 is controlled by the a_2 coefficient appearing in (2) and the result is:

$$\langle \sigma \pi^0 | H_{\text{eff}} | \bar{B}^0 \rangle = \frac{G_F}{\sqrt{2}} V_{ub}^* V_{ud} a_2 F_0^{(B\sigma)} (m_\pi^2) (m_B^2 - m_\sigma^2) f_\pi \times \frac{g_{\sigma\pi^+\pi^-}}{u - m_\sigma^2 + i\Gamma_\sigma(u)m_\sigma}, \quad (5)$$

with $a_2 = C_2 + \frac{1}{3}C_1$. We have checked that the formulas (3) and (5), when applied to $D \rightarrow \sigma\pi$ with the obvious changes in the masses, form factor values and CKM coefficients, reproduce the experimental results of [15].

The definition of the form factors $F_0^{(B\sigma)}$ and $F_1^{(B\sigma)}$ is the following:

$$\langle \sigma(q_\sigma) | A^\mu(q) | B(p) \rangle = \left[\frac{m_D^2 - m_\sigma^2}{q^2} q^\mu \right] F_0^{(B\sigma)}(q^2) + \left[(p + q_\sigma)^\mu - \frac{m_D^2 - m_\sigma^2}{q^2} q^\mu \right] F_1^{(B\sigma)}(q^2), \quad (6)$$

with $F_1(0) = F_0(0)$. Applying the factorization selects the form factor $F_0^{(B\sigma)}$ since:

$$\langle \pi^- | A_{(du)}^\mu | \text{VAC} \rangle = i f_\pi q^\mu. \quad (7)$$

The *polar* and *direct* contributions to the semileptonic form factors discussed in [17] for the $D \rightarrow \sigma$ transition, give in this case:

$$F_0^{(B\sigma)}(m_\pi^2) \simeq F_0(0) = 0.45 \pm 0.15, \quad (8)$$

where $m_B = 5.27$ GeV, $m_B(1^+) = 5.7$ GeV, $B(1^+)$ being the polar intermediate state connecting B to A^μ [17].

The errors include the uncertainty due to the variation of the parameters of the CQM model [18] in a fixed range of values and that arising from the extrapolation of the polar form factors to $q^2 = m_\pi^2 \simeq 0$, following the same steps as in [17]. In that paper the analogous form factor for D decays was evaluated, $F_0^{(D\sigma)}$, and the result of the model closely reproduces the experimental value of [15]. The calculation does not include $1/m_b$ corrections, where m_b is the heavy quark mass, however such corrections were estimated to be smaller than the quoted experimental error for D decays and are even smaller in the study of B decays.

The possibility of a σ contribution to CP violations was studied in $K \rightarrow \pi\pi$ decays [19]. We consider in the following the impact of σ contribution in quasi two-body pion decays of the B meson. In particular the study of the $B \rightarrow \rho\pi$ channels is used for the determination of the unitarity angles α and γ [4]. These analyses make the assumption that, using cuts in the three invariant masses for the pion pairs, one can extract the ρ contribution without significant background contaminations. The ρ has spin 1, the π spin 0 as well as the initial B , and therefore the ρ has angular distribution $\cos^2\theta$ (θ is the angle of one of the ρ decay products with the other π in the ρ rest frame). This means that the Dalitz plot is mainly populated at the border, especially the corners, by this decay. Analyses following these lines were performed by the BABAR working groups [2]; MonteCarlo simulations, including the background from the narrow f_0 resonance, show that, with cuts at $m_{\pi\pi} = m_\rho \pm 200$ MeV, no significant contributions from other sources are obtained. The role of excited resonances such as the ρ' and the non-resonant background was discussed [20]. Finally the role of the off mass-shell B^* contribution was discussed in [9,11,12].

The formulas derived in the present paper allow to estimate the contribution of the σ which was not taken into account in the previous analyses. As the resonance is broad, part of the events from $B \rightarrow \sigma\pi \rightarrow 3\pi$ survive the cuts on the invariant mass of two π that reconstruct the mass of the ρ (within ± 200 MeV) in $B \rightarrow \rho\pi \rightarrow 3\pi$. We define the integrating region in the Dalitz plot around the ρ resonance:

$$\Gamma_{eff}(\bar{B}^0 \rightarrow \rho^- \pi^+) = \Gamma(\bar{B}^0 \rightarrow \pi^+ \pi^- \pi^0) \Big|_{m_\rho - \delta \leq \sqrt{s} \leq m_\rho + \delta}$$

$$\Gamma_{eff}(\bar{B}^0 \rightarrow \rho^+ \pi^-) = \Gamma(\bar{B}^0 \rightarrow \pi^+ \pi^- \pi^0) \Big|_{m_\rho - \delta \leq \sqrt{t} \leq m_\rho + \delta}.$$

The Mandelstam variables are $s = (p_{\pi^+} + p_{\pi^0})^2$, $t = (p_{\pi^-} + p_{\pi^0})^2$ and we use $\delta = 200$ MeV.

Moreover the σ contributes to the decay $B^\pm \rightarrow \pi^+ \pi^- \pi^\pm$ and only in a negligible way to the decay $\bar{B}^0 \rightarrow \pi^+ \pi^- \pi^0$ if its contribution is restricted to the experimental cuts that reconstruct the process $\bar{B}^0 \rightarrow \rho^\pm \pi^\mp$. Note that in the latter case the ρ meson is charged, while the σ

is neutral. This means that on average the two charged pions will have a high invariant mass from the σ resonance in the process, while in order to reconstruct a charged ρ meson, a neutral and a charged pion have to be used. We have checked numerically that this is indeed correct and the σ contribution to $\bar{B}^0 \rightarrow \rho^\pm \pi^\mp$ (we reconstruct the ρ mass within ± 200 MeV) is three orders of magnitude smaller than the ρ contribution (for which we find $\mathcal{B}(\bar{B}^0 \rightarrow \rho^\mp \pi^\pm) = 19.9 \times 10^{-6}$). The σ contribution to the charged B decay is on the contrary a fraction of the ρ one and comparable or larger than the contribution from the B^* (see Table II). This provides a clear mechanism to enhance the denominator of the ratio R , giving a result closer to the experimental one:

$$R = \frac{\mathcal{B}(\bar{B}^0 \rightarrow \rho^\mp \pi^\pm)}{\mathcal{B}(B^- \rightarrow \rho^0 \pi^-)} = 3.6, \quad (9)$$

including only the ρ and σ contributions, or $R \simeq 4$ including also the B^* , which however is less precisely known. A possibility to disentangle these contributions from the ρ is to vary the experimental cuts and see the effect on the effective branching ratios. We find that the process $B^- \rightarrow \sigma \pi^-$ has a total branching ratio of 4.3×10^{-6} , therefore comparable to the one of the ρ . Allowing the cuts around the ρ mass to be ± 300 MeV, the σ contribution grows to 2.7×10^{-6} , to be compared to what indicated in Table II for a cut of ± 200 MeV.

ACKNOWLEDGMENTS

ADP acknowledges support from EU-TMR programme, contract CT98-0169. We wish to thank N. A. Törnqvist, R. Gatto and G. Nardulli for informative discussions.

-
- [1] Y. Kubota et al., Nucl. Inst. and Meth. A320, 66 (1992).
[2] P. F. Harrison and H. R. Quinn [BABAR Collaboration], SLAC-R-0504 *Papers from Workshop on Physics at an Asymmetric B Factory (BaBar Collaboration Meeting)*, <http://www.slac.stanford.edu/pubs/slacreports/slac-r-504.html>.
[3] BELLE Technical Design Report, KEK-Report 95-1, April 1995, <http://bsunsrv1.kek.jp/bdocs/tdr.html>; F. Takasaki, 19th International Symposium on Lepton and Photon Interactions at High Energies (LP99), Stanford, California, 9-14 Aug. 1999 hep-ex/9912004.
[4] see for example H. J. Lipkin, Y. Nir, H. R. Quinn and A. Snyder, Phys. Rev. **D44**, 1454 (1991); A. E. Snyder and H. R. Quinn, Phys. Rev. **D48**, 2139 (1993); G. Eilam, M. Gronau and R. R. Mendel, Phys. Rev. Lett. **74**, 4984 (1995) [hep-ph/9502293]; I. Bediaga, R. E. Blanco, C. Gobel and R. Mendez-Galain, Phys. Rev. Lett. **81**, 4067 (1998) [hep-ph/9804222]; R. E. Blanco, C. Gobel and R. Mendez-Galain, hep-ph/0007105.
[5] C. P. Jessop et al. [CLEO Collaboration], hep-ex/0006008.
[6] B. Aubert et al. [BABAR Collaboration], SLAC-PUB-8537.
[7] Y. Gao and F. Wurthwein [CLEO Collaboration], hep-ex/9904008.
[8] M. Bauer, B. Stech and M. Wirbel, Z. Phys. **C34**, 103 (1987).
[9] A. Deandrea, R. Gatto, M. Ladisa, G. Nardulli and P. Santorelli, Phys. Rev. **D62**, 036001 (2000) [hep-ph/0002038];
[10] M. Ciuchini, E. Franco, G. Martinelli and L. Silvestrini, Nucl. Phys. **B501**, 271 (1997) [hep-ph/9703353]; M. Ciuchini, R. Contino, E. Franco, G. Martinelli and L. Silvestrini, Nucl. Phys. **B512**, 3 (1998); Erratum-ibid. **B531**, 656 (1998) [hep-ph/9708222].
[11] A. Deandrea, proceedings of 35th Rencontres de Moriond: Electroweak Interactions and Unified Theories, Les Arcs, France, 11-18 Mar 2000, hep-ph/0005014;
[12] A. Deandrea, R. Gatto, M. Ladisa, G. Nardulli and P. Santorelli, hep-ph/0007059.
[13] N. G. Deshpande, G. Eilam, X. He and J. Trampetic, Phys. Rev. **D52**, 5354 (1995) [hep-ph/9503273];
[14] D. E. Groom et al., Eur. Phys. J. **C15**, 1 (2000); N. A. Tornqvist and M. Roos, Phys. Rev. Lett. **76**, 1575 (1996) [hep-ph/9511210].
[15] E. M. Aitala et al. [E791 Collaboration], hep-ex/0007028; J. Slaughter, talk at XXXVth Rencontres de Moriond on Hadronic interactions and QCD, <http://moriond.in2p3.fr/QCD00/transparencies/>
[16] A. D. Polosa, N. A. Tornqvist, M. D. Scadron and V. Elias, hep-ph/0005106; C. Dib and R. Rosenfeld, hep-ph/0006145.
[17] R. Gatto, G. Nardulli, A. D. Polosa and N. A. Tornqvist, hep-ph/0007207.
[18] A. Deandrea, N. Di Bartolomeo, R. Gatto, G. Nardulli and A. D. Polosa, Phys. Rev. **D58**, 034004 (1998) [hep-ph/9802308]; *ibid.* **D59**, 074012 (1999) [hep-ph/9811259]; *ibid.* **D61**, 017502 (2000) [hep-ph/9907225]; A. Deandrea, R. Gatto, G. Nardulli and A. D. Polosa, JHEP **9902**, 021 (1999) [hep-ph/9901266]; A. D. Polosa, hep-ph/0004183.
[19] T. Morozumi, C. S. Lim and A. I. Sanda, Phys. Rev. Lett. **65**, 404 (1990); Y. Keum, U. Nierste and A. I. Sanda, Phys. Lett. **B457**, 157 (1999) [hep-ph/9903230]; K. Terasaki, hep-ph/0008225.
[20] J. Charles, PhD Thesis (in French); S. Versille, PhD Thesis (in French).

TABLE I. Estimates for the ratio R beyond the factorization approximation (the so-called charming penguins) using different sets of input data: QCD sum-rules (QCDSR), lattice-QCD, quark models (QM)

QCDSR	$R = 6.3$
lattice	$R = 5.5$

TABLE II. Effective branching ratios (in units of 10^{-6}) for the charged $B^- \rightarrow \pi^+\pi^-\pi^-$ decay into three pions. Cuts as indicated in the text. Set I refers to the choice of hadronic parameter $g = 0.4$ for the coupling $BB^*\pi$ relevant for the evaluation of the B^* contribution, while Set II is calculated using $g = 0.6$. The σ contribution is less dependent on hadronic uncertainties as the coupling $g_{\sigma\pi^+\pi^-}$ is obtained from data.

$B^- \rightarrow \pi^+\pi^-\pi^-$	ρ	σ	B^*	$\rho + \sigma$	all
Set I	3.8	1.5	0.8	5.5	4.9
Set II	3.8	1.5	1.8	5.5	5.1




FIG. 1. Tree level diagram for a B^- decaying to three pions via the σ resonance. The two identical π^- in the final state require to sum coherently two such diagrams having the momentum labels p_1 and p_2 inverted.




FIG. 2. As in Fig. 1 but for the $\bar{B}^0$, with no identical particles in the final state.